

Water footprint and life cycle assessment of *Populus spp.* bioenergy system: a case study in Southern Europe

E. Seigné-Itoiz^{1,*}, C.M. Gasol^{1,2}, F. Brun³, L.Rovira⁴, J.M Pagés⁴, F.Camps⁴, J.Oliver^{1,2}, J. Rieradevall^{1,5}, X. Gabarrell^{1,5}

¹Institute of Environmental Science and Technology (ICTA). SosteniPrA (ICTA-IRTA-Inèdit). Universitat Autònoma de Barcelona (UAB). 08193 Bellaterra (Barcelona, Spain)

²Inèdit Innovació SL. SosteniPrA (ICTA-IRTA-Inèdit). UAB Research Park. 08193 Bellaterra (Barcelona, Spain)

³Dipartimento di Economia e Ingegneria Agraria Forestale e Ambientale, University of Torino (Italy).

⁴IRTA-Mas Badia. Institut de Recerca i Tecnologia Agroalimentaria (IRTA). Estación Experimental Fundación Mas Badia Ctra. De la Tallada, s/n 17134 La Tallada (Girona)

⁵Chemical Engineering Department, XRB. Universitat Autònoma de Barcelona (UAB). 08193 Bellaterra (Barcelona, Spain)

*Corresponding author: eva.sevine@campus.uab.cat Tel.: (+34) 935 813 760 - Fax.: (+34) 935 813 331

1 Introduction

Short rotation energy crops have been promoted because they provide biomass in short periods of time but the impacts of water consumption on existing water resources have not been analyzed in depth. Previous studies has conducted of *Populus spp.* bioenergy systems^{1,2} focus on environmental impacts and it has detected that water consumption will be crucial for Southern Europe. This study evaluates the **water footprint (WF) and the life cycle assessment (LCA) of a *Populus spp.* crop in Spain through its first cycle for two planting densities in order to evaluate if water consumption would be different for the same energy production.**



- ✓ Low input agrochemical
- ✓ Cycle: two years
- ✓ Low Density: 6,666 pl ha⁻¹
- ✓ High Density: 20,000 pl ha⁻¹

High density
Crop of *Populus spp.* bioenergy system studied

3 Methodology

For the water consumption has been used the methodology of water footprint (WF)^{3,4} and for the energy and environmental assessment it has been used the life cycle analysis methodology. Grey WF has rejected due to low input of agrochemical requirements.

3.1 Life cycle assessment- Functional Unit

To obtain 1 ha of *Populus spp.* destined to produce biomass throughout the first cycle of *Populus spp.* bioenergy system.

3.3 Life cycle Inventory and life cycle impact assessment


Data has been collected throughout the first cycle of the crop and ecoinvent 2.0 (2007) databases. The LCIA methodology used has been "Cumulative Energy Demand v 1.4" and "CML baseline 2001".

	Low Density		High Density	
	Indirect water	Direct water	Indirect water	Direct water
Agrochemical production (m ³ ha ⁻¹ yr ⁻¹)	310		507	
Agricultural machinery (m ³ ha ⁻¹ yr ⁻¹)	24		269	
Diesel (m ³ ha ⁻¹ yr ⁻¹)	12		21	
Irrigation (m ³ ha ⁻¹ yr ⁻¹)	-	1,808	-	3,288

Water life cycle inventory of *Populus spp.* bioenergy system for LD and HD


2 Systems description

The system under study includes all stages of agricultural production for the first cycle of the crop: work in the plantation, production of agrochemicals, tractor and agricultural utensils manufacture as water consumption through life cycle.



4 Results and discussion


Biomass and energy per ha obtained; and blue WF and green WF for both planting densities per ton and per GJ obtained and the biomass and energy obtained.



	LD (m ³ ton ⁻¹)	HD (m ³ ton ⁻¹)
Blue WF	266	396
Green WF	754	628
Total WF	1020	1023

	LD (m ³ GJ ⁻¹)	HD (m ³ GJ ⁻¹)
Blue WF	7	11
Green WF	21	17
Total WF	28	28

Figures present the relative contributions of life cycle stages of low density and high density for the first cycle of *Populus spp.*, respectively.



Relative contributions of life cycle stages of LD

Relative contributions of life cycle stages of HD

5 Conclusions

The total WF is high for bioenergy production of *Populus spp.* in Southern Europe and per ton produced and for GJ obtained, values are similar for both densities consider (28m³ · GJ⁻¹). However for high density a higher amount of water is required from the ground or surface area and this type of planting density could increase water demand in the area(3,288 m³ ·ha⁻¹ · yr⁻¹). Bioenergetic system of *Populus spp.* in southern Europe should be restricted to areas where water availability is abundant and low density design is preferred to higher densities. Results from LCA indicated that for both densities the highest environmental impacts are due to agricultural machinery production. Low input design in agrochemical consumption implies a reduction of environmental impacts.

References

- ¹Gasol C.M., Gabarrell X., Anton A., Rigola M., Carrasco J., Ciria P. & Rieradevall J. LCA of populus spp. bioenergy system compared with Brassica carinata energy crop and natural gas in regional scenario. Biomass and Bioenergy 33 (2009), 119-129
- ²Gerbens-Leenes, W., Hoekstra, A.Y. and Van der Meer, T.H. (2009) The water footprint of bioenergy. Proceedings of the National Academy of Sciences 106 (25): 10219-10223.
- ³Hoekstra, A.Y. and Chapagain, A.K. (2007) Water footprints of nations: water use by people as a function of their consumption pattern, Water Resources Management 21(1): 35-48.
- ⁴Hoekstra, A. Y., Chapagain, A. K., Aldaya, M. M. and Mekonnen, M. M. (2009) Water footprint manual: State of the art 2009, Water Footprint Network, Enschede, the Netherlands, www.waterfootprint.org/downloads/WaterFootprintManual2009.pdf (retrieved 15 February 2010).